



UNIVERSITÄT HOHENHEIM



# Phenological Responses of Upland Rice Grown Along an Altitude Gradient

Suchit Shrestha<sup>1</sup>, Folkard Asch<sup>1</sup>, Holger Brück<sup>1</sup>, Alain Ramanantsoanirina<sup>2</sup>, Julie Dusserre<sup>3</sup>

<sup>1</sup>University of Hohenheim, Germany; <sup>2</sup>FOFIFA, Madagascar and <sup>3</sup>CIRAD, France

International conference on crop improvement, ideotyping and modeling  
for African cropping systems under climate change  
(CIMAC)



Developing Rice and SOrgnum Crop Adaptation Strategies for climate change in vulnerable environments in Africa

7 - 9 February 2011, Stuttgart-Hohenheim, Germany.



## INTRODUCTION

- Crop adaptation strategies are required for varietal development and crop management to avoid negative impacts of climate change.
- Characterization of potential upland rice ideotype traits is necessary to fit genotype into the crop production system.
- Phenology of upland rice responds to temperature/altitude gradient.
- Spikelet sterility needs to be considered along with crop phenology to fit genotype to the crop production system.



## **OBJECTIVES**

- To identify phenological responses to altitude gradients.
- To assess the impact of sowing date on spikelet sterility in contrasting environments.




## MATERIALS and METHODS

### Locations

High altitude (Andranomanelatra)

Mid altitude (Ivory)

Low altitude (Ankepaka)



|                      | Location       |                |                |
|----------------------|----------------|----------------|----------------|
|                      | High altitude  | Mid altitude   | Low altitude   |
| <b>Latitude</b>      | 19°46'45.3" S  | 19°33'16.8" S  | 22°11'31.6" S  |
| <b>Longitude</b>     | 47°06'24.5" E  | 46°25'29.1" E  | 47°52'32.7" E  |
| <b>Elevation</b>     | 1625 m asl     | 965 m asl      | 25 m asl       |
| <b>Altitude</b>      | High           | Mid            | Low            |
| <b>Precipitation</b> | 1300 mm        | 1200 mm        | 2100 mm        |
| <b>Tmin - Tmax</b>   | 12.6 – 18.9 °C | 19.4 – 23.8 °C | 19.2 – 27.1 °C |
| <b>Soil texture</b>  | Clay           | Clay loam      | Silt loam      |




## Meteorology

| <u>Sowing</u> | <u>High altitude</u> |
|---------------|----------------------|
| 1             | Mid Sep              |
| 2             | Mid Oct              |
| 3             | Mid Nov              |
| 4             | Mid Dec              |
| 5             | Mid Jan              |

| <u>Sowing</u> | <u>Mid altitude</u> |
|---------------|---------------------|
| 1             | Mid Sep             |
| 2             | Mid Oct             |
| 3             | Mid Nov             |
| 4             | Mid Dec             |
| 5             | Mid Jan             |

| <u>Sowing</u> | <u>Low altitude</u> |
|---------------|---------------------|
| 1             | Mid Oct             |
| 2             | Mid Nov             |
| 3             | Mid Dec             |
| 4             | Mid Jan             |
| 5             | Mid Feb             |







## Genotypes

| Genotypes    | Sub-species  | Specific traits     | Growing altitude |
|--------------|--|---------------------|------------------|
| B 22         | <i>O. Sativa</i> (tropical japonica)                                     | Plasticity (aspect) | Mid-Low          |
| Botramaintso | <i>O. Sativa</i> (tropical japonica)                                     | Vigor growth        | Mid              |
| Chomrong     | <i>O. Sativa</i> (temperate japonica)                                    | Cold tolerant       | High             |
| FOFIFA 161   | <i>O. Sativa</i> (tropical japonica)                                     | Cold tolerant       | High             |
| FOFIFA 167   | <i>O. Sativa</i> (tropical japonica)                                     | Cold tolerant       | High             |
| FOFIFA 172   | <i>O. Sativa</i> (tropical japonica)                                     | Cold tolerant       | High             |
| IRAT 112     | <i>O. Sativa</i> (tropical japonica)                                     | Plasticity (aspect) | Mid              |
| Nerica 4     | <i>O. Glaberrima</i> x <i>O. Sativa</i> (interspecific crosses)          | Stay green          | Mid              |
| Primavera    | <i>O. Sativa</i> (tropical japonica)                                     | Grain quality       | Mid-Low          |
| WAB 878      | <i>O. Glaberrima</i> x <i>O. Sativa japonica</i> (interspecific crosses) | Vigor growth        | Mid              |

|         | B22 | Botramaintso | Chhomrong | FOFIFA 161 | FOFIFA 167 | FOFIFA 172 | IRAT 112 | NERICA 4 | PRIMAVERA | WAB 878 |
|---------|-----|--------------|-----------|------------|------------|------------|----------|----------|-----------|---------|
| Tiller  |     |              |           |            |            |            |          |          |           |         |
| Canopy  |     |              |           |            |            |            |          |          |           |         |
| Panicle |     |              |           |            |            |            |          |          |           |         |
| Grain   |     |              |           |            |            |            |          |          |           |         |



- 10 varieties of rainfed upland rice were planted at 5 monthly staggered sowing dates without replication in three locations for two years (phenological garden).
- Plot size was 1 X 1 m with 20 X 20 cm spacing between plants.
- Early sowing plots were manually irrigated to avoid drought stress.
- Phenological stages were carefully observed during the crop cycle.
- Biomass, yield, yield components and sterility percentage were determined at maturity.





# Phenological Garden (Mini Rice Garden)








## RESULTS and DISCUSSION


### Calendar days



High altitude conditions increase crop duration by 36 to 52 days.




## Thermal days (10°C base temperature used for all varieties)



The local landrace has higher thermal demand.



## Sowing dates and crop duration (all varieties photoperiod insensitive)




Crop duration is only defined by temperature (via altitude).



## Altitudes alter crop duration

| Genotype     | Slope | Intercept | r <sup>2</sup> |
|--------------|-------|-----------|----------------|
| B22          | -6.5  | 239       | 0.800          |
| Botramaintso | -8.9  | 314       | 0.851          |
| Chhomrong    | -6.3  | 236       | 0.807          |
| FOFIFA 161   | -5.9  | 225       | 0.823          |
| FOFIFA 167   | -6.9  | 254       | 0.778          |
| FOFIFA 172   | -4.8  | 199       | 0.554          |
| IRAT 112     | -6.2  | 231       | 0.832          |
| NERICA 4     | -6.0  | 226       | 0.823          |
| Primavera    | -6.5  | 244       | 0.851          |
| WAB 878      | -7.2  | 258       | 0.862          |

Rise in air temperature by 1°C decreases crop duration by 5 to 9 days from germination to flowering depending on genotype.








## Spikelet sterility in different altitudes

Low temperature effects on sterility

Cold tolerant varieties perform well at high altitude

Genetic adaptation to cold environments can be exploited





## **CONCLUSION and OUTLOOK**

- Predicted rise in air temperature is favourable for upland rice cultivation in the high altitude in terms of crop duration and grain yield.
- Genetic variation in cold tolerance can be used to adapt rice to cold environments.
- Morpho-physiological traits contributing to cold tolerance need to be identified for further breeding.



## ACKNOWLEDGEMENT



Bundesministerium für  
wirtschaftliche Zusammenarbeit  
und Entwicklung

**Funding organization**  
**German Ministry for Collaboration and**  
**Development through GIZ/BEAF.**



**Africa Rice Center**

**Collaborator**



**Collaborator**



**Co-partner institute**



UNIVERSITÄT HOHENHEIM



# THANK YOU



Developing Rice and Sorghum Crop Adaptation Strategies for climate change in vulnerable environments in Africa

7 - 9 February 2011, Stuttgart-Hohenheim, Germany.